BK030062

Section 8-510(k) Summary of Safety and Effectiveness

8.1 Statement This summary of 510(k) safety and effectiveness information is being submitted

in accordance with the requirements of SMDA 1990 and CFR 807.92

8.2

Submitter Haem

Haemonetics Corporation

400 Wood Road

Braintree, MA. 02184-9114

8.3

Susan Finneran

Company Contact RA Product Manager 355 Wood Road

Braintree, MA.

781-356-9220

8.4

Proprietary Name: Haemonetics® Model 215 Automated Glycerolization /

Device

Deglycerolization System Common Name: ACP 215

Name

Classification Name: Processing System for Frozen Blood

8.5

Predicate

Legally

Marketed

Devices

The Haemonetics ACP 215, which is the subject of this submission, is substantially equivalent to the previously cleared ACP 215 cleared via

BK000002 on May 4, 2001 by Haemonetics Corporation.

8.6

Device

Description

The ACP 215 is an automated cell processing system intended to be used glycerolize and red blood cells.

8.7
Device
Indications
and Intended
use

Indications for use:

The ACP 215 is intended to glycerolize and deglycerolize red blood cells derived from whole blood that have been stored in any approved anticoagulant/additive solution for 6 days at 1-6°C prior to glycerolization. Red blood cells that have been collected and stored in CPDA-1, CPD/AS-1, or CP2D/AS-3, glycerolized using the ACP 215, frozen at -65 °C or colder, deglycerolized by the ACP 215, and stored in Haemonetics AS-3 at 1-6° may be labeled for extended storage for 14 days.

Red blood cells collected in any other anticoagulant / additive solution will be labeled with a 24-hour outdate.

The following page contains a table comparing the new versus the predicate device.

Characteristics	SUSTANTIAL EQUIVALENCE Predicate Device	Proposed Device
Character istics	ACP 215 System	ACP 215 System
Protocol	Automated Glycerolization/ Deglycerolization	Automated Glycerolization/ Deglycerolization
Hardware	ACP 215	ACP 215
Disposable	LN 225-00 and LN235-00	LN 225-00 and LN235-00 (or LN236-00)
Washing	12 %NaCl	12 %NaCl
Solution	0.9% NaCl/ 0.2gm Glucose	0.9% NaCl/ 0.2gm Glucose
Intended Use	The ACP 215 is intended to Glycerolize and Deglycycerolize Red blood Cells up to 6 days old. Cells collected and stored in CPDA-1 may be labeled for extended storage up to 14 days. Cell collected in other solutions may be stored for 24 hours post-deglycerolization.	The ACP 215 is intended to glycerolize and deglycerolize red blood cells derived from whole blood that have been stored in any approved anticoagulant/additive solution for 6 days at 1-6°C prior to glycerolization. Red blood cells that have been collected and stored in CPDA-1, CPD/AS-1, or CP2D/AS-3, glycerolized using the ACP 215, and frozen at -65 °C or colder, deglycerolized by the ACP 215, and stored in Haemonetics AS-3 at 1-6° may be labeled for extended storage for 14 days. Red blood cells collected in any other anticoagulan / additive solution will be labeled with a 24-hour outdate.
Product Quality	Recovery: 95% confidence that 90% units met the 80% minimum recovery Hemolysis: 95% confident that at 90% of the units met the maximum hemolysis of 1%	Recovery: 95% confidence that 90% units met the 80% minimum recovery Hemolysis: 95% confident that at 95% of the units met the maximum hemolysis of 1%
Post-Washing Storage	14-days post deglycerolization storage.	14-days post deglycerolization storage